

F



Dedicate®

DEDICATE® is a contact and systemic fungicide which has early curative and preventative action against Fusarium Patch (*Microdochium nivale*), Red Thread (*Laetisaria fuciformis*), Dollar Spot (*Sclerotinia homoeocarpa*), Anthracnose (*Colletotrichum cereale*) Leaf Spot (*Drechslera* spp.) and Rust (*Puccinia* spp) in managed amenity turf, at any time of year.

Fungicide

MAPP 13612

A suspension concentrate formulation containing
200 g/l (18.2% w/w) tebuconazole and
100 g/l (9.10% w/w) trifloxystrobin

250 ml e



HARMFUL

VERY TOXIC TO AQUATIC ORGANISMS, MAY CAUSE LONG-TERM ADVERSE EFFECTS IN THE AQUATIC ENVIRONMENT. POSSIBLE RISK OF HARM TO THE UNBORN CHILD. IRRITATING TO THE RESPIRATORY SYSTEM.

Do not breathe spray.

This material and its container must be disposed of in a safe way.

Wear suitable protective clothing and gloves.

Use appropriate containment to avoid environmental contamination.

May produce an allergic reaction. Contains trifloxystrobin.

To avoid risks to man and the environment comply with the instructions for use.

DEDICATE®

Containing 200 g/l (18.2% w/w) tebuconazole and
100 g/l (9.10% w/w) trifloxystrobin



DANGEROUS FOR THE ENVIRONMENT

For important information and detailed directions for use please see inside leaflet.

For 24 hour emergency information contact Bayer CropScience Ltd
Telephone: 0800 220876 or nearest National Poisons Information Centre.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

WHO Classification: III (slightly hazardous)



The
Voluntary
Initiative

Batch number : see on the pack

This label is compliant with the CPA
Voluntary Initiative Guidance



Bayer Environmental Science

A Business Operation of Bayer CropScience

Bayer Environmental Science
Bayer CropScience Limited
230 Cambridge Science Park
Milton Road
Cambridge CB4 0WB
Tel: 00800 1214 9451

Important Information

FOR USE ONLY AS A HORTICULTURAL FUNGICIDE.

For use on managed amenity turf and amenity grassland.

Maximum individual dose: 1 L/ha

Maximum number of treatments: 2 per year

Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence.

Follow the code of practice for using plant protection products.

SAFETY PRECAUTIONS**Operator Protection**

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment :

WEAR SUITABLE PROTECTIVE CLOTHING AND SUITABLE PROTECTIVE GLOVES when handling the concentrate, applying by hand-held equipment or handling contaminated surfaces. However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

DO NOT BREATHE SPRAY.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH ANY CONTAMINATION from eyes immediately.

IF YOU FEEL UNWELL, seek medical advice (show label where possible).

WASH HANDS AND EXPOSED SKIN before meals and after work.

Environmental Protection

DO NOT CONTAMINATE SURFACE WATERS OR DITCHES with chemical or used container.

Storage and Disposal

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER tightly closed in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Follow the code of practice for using plant protection products.

DIRECTIONS FOR USE:

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

For use on established turf. DEDICATE® may be applied to turf sites including golf courses, institutional, commercial and residential lawns, sports fields, parks, municipal grounds and cemeteries.

GENERAL INFORMATION:

DEDICATE® is a fungicide exhibiting both contact and systemic mode of action. It is a broad spectrum triazole and strobilurin fungicide recommended for control or reduction of a range of diseases on managed amenity turf.

RESTRICTIONS:

Do not apply during drought conditions or to frozen turf.

APPLICATION:

For control of Fusarium, Red Thread and Dollar Spot, use 1 L of DEDICATE® in a minimum of 400 - 500 litres of water per hectare of turf. Control of Leaf Spot and reduction of Rust and Anthracnose may also be achieved. For use with knapsack sprayers or vehicle mounted hydraulic sprayers.

Application Equipment			
Vehicle-mounted Hydraulic Sprayer		Knapsack Sprayer	
Rate per hectare	Water per hectare	Rate per 100 m ²	Water per 100 m ²
1 Litre	400-500 litres	10 mls	4 -5 litres
		Rate per 500 m ²	Water per 500 m ²
		50 mls	20-25 litres

Application Timing

DEDICATE® should be used preventatively or as early as possible in the disease cycle, applying at the first sign of disease.

DEDICATE® can be used when grass is either dormant or activity actively growing and displays both protective and some curative activity. If grass is being mown, DEDICATE® should be applied after cutting and further mowing should not take place until AT LEAST

48 hours after treatment. This is to allow adequate movement of the systemic active ingredients throughout the plant.

Do not apply more than 2 applications per year.

Allow at least 28 days between applications of DEDICATE®.

Do not apply sequential applications of DEDICATE®.

Alternate with a fungicide having a different mode of action.

Resistance Management:

DEDICATE® contains trifloxystrobin which belongs to the strobilurin class of chemistry and also contains Tebuconazole, of the Dimethylation class of chemistry and two separate classes of fungicide with different modes of action together in one product should reduce the chance of resistance appearing however the possible development of diseases resistant to DEDICATE® cannot be excluded or predicted. Certain fungal pathogens are known to develop resistance to strobilurin products when used repeatedly. The use of DEDICATE® should conform to resistance management strategies. Such strategies may include rotating with products containing active substances with different modes of action, or limiting the total number of applications per season. Bayer recommends responsible product stewardship to ensure effective long term control of the fungal diseases on this label.

Mixing:

Sprayers should be THOROUGHLY CLEANED before use and filters and jets checked for damage and blockages.

Thoroughly shake the pack before use.


Half fill the sprayer tank with clean water and agitate. Add the required amount of DEDICATE®.


Mix thoroughly and make up to the required volume of water while still agitating. Spray immediately after mixing. Continue agitation at all times during spraying and stoppages, until the tank is completely empty.


Wash out all spray equipment thoroughly with water immediately after use.


DEDICATE® is a registered trademark of Bayer Environmental Science SAS


Color : BLACK





























F



Dedicate®

DEDICATE® is a contact and systemic fungicide which has early curative and preventative action against Fusarium Patch (*Microdochium nivale*), Red Thread (*Laetisaria fuciformis*), Dollar Spot (*Sclerotinia homoeocarpa*), Anthracnose (*Colletotrichum cereale*) Leaf Spot (*Drechslera* spp.) and Rust (*Puccinia* spp) in managed amenity turf, at any time of year.

Fungicide

MAPP 13612

A suspension concentrate formulation containing 200 g/l (18.2% w/w) tebuconazole and 100 g/l (9.10% w/w) trifloxystrobin

250 ml e



HARMFUL

VERY TOXIC TO AQUATIC ORGANISMS, MAY CAUSE LONG-TERM ADVERSE EFFECTS IN THE AQUATIC ENVIRONMENT. POSSIBLE RISK OF HARM TO THE UNBORN CHILD. IRRITATING TO THE RESPIRATORY SYSTEM.

Do not breathe spray.

This material and its container must be disposed of in a safe way.

Wear suitable protective clothing and gloves.

Use appropriate containment to avoid environmental contamination.

May produce an allergic reaction. Contains trifloxystrobin.

To avoid risks to man and the environment comply with the instructions for use.

DEDICATE®

Containing 200 g/l (18.2% w/w) tebuconazole and 100 g/l (9.10% w/w) trifloxystrobin



DANGEROUS FOR THE ENVIRONMENT

For important information and detailed directions for use please see inside leaflet.

For 24 hour emergency information contact Bayer CropScience Ltd

Telephone: 0800 220876 or nearest National Poisons Information Centre.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

WHO Classification: III (slightly hazardous)



The Voluntary Initiative

Batch number : see on the pack

This label is compliant with the CPA Voluntary Initiative Guidance



Bayer Environmental Science

A Business Operation of Bayer CropScience

Bayer Environmental Science
Bayer CropScience Limited
230 Cambridge Science Park
Milton Road
Cambridge CB4 0WB
Tel: 00800 1214 9451